





Primary* Refugee Arrivals to MN by Region of World 1979-2015







Primary Refugee Arrival by Month, Minnesota, 2011-2015



2015 Primary Refugee Arrivals To Minnesota (N=2,244)




Number of Refugee Arrivals By Initial County Of Resettlement









Primary Refugee Arrivals, Minnesota 2015




“Other” includes Afghanistan, Belarus, Cameroon, Rep. of Congo, Cuba, DR Congo, Eritrea, Honduras, Iran, Liberia, Mexico, Moldova, Nepal, Russia, Sri Lanka, Sudan, Syria, Tanzania, Ukraine, and Vietnam



Country of Origin by County of Resettlement, 2015




Primary Refugee Arrivals Screened Minnesota, 2005-2015*



Ineligible if moved out of state or to an unknown destination, no insurance, unable to locate or died before screening

Primary Refugees Lost to Follow-up Minnesota, 2015






Primary Refugee Screenings by Region of Origin, Minnesota, 2015

World Region	Total arrivals	Ineligible for Screening	Number Screened (%*)
Sub-Saharan Africa	1,150	28	1,109 (99%)
SE Asia/E Asia	884	4	880 (100%)
North Africa/ Middle East	149	0	122 (82%)
Eastern Europe	58	1	57 (98%)
Latin America/ Caribbean	3	2	1 (100%)

Refugee Screening Rates by Exam Type Minnesota, 2015





Health Status of New Refugees, Minnesota, 2015*

Health status upon arrival	No. of refugees	No. (%) with infection
TB (latent or active)**	2,137 (97%)	425 (20%)
Hepatitis B infection***	2,084 (98%)	87 (4%)
Parasitic Infection****	1,909 (88%)	326 (17%)
Sexually Transmitted Infections (STIs)*****	2,095 (97%)	17 (1%)
Malaria Infection	96 (4%)	0 (0%)
Lead*****	949 (96%)	64 (7%)
Hemoglobin	2,130 (98%)	478 (23%)

*Total screened: **N=2,168** (98% of 2,209 eligible refugees)

** Persons with LTBI (≥ 10 mm induration or IGRA+, normal CXR) or suspect/active TB disease

*** Positive for Hepatitis B surface antigen (HBsAg)

**** Positive for at least one intestinal parasite infection


***** Positive for at least one STI (tested for syphilis, HIV, chlamydia or gonorrhea)

***** Children <17 years old (N=992 screened); lead level ≥ 5 $\mu\text{g}/\text{dL}$



Tuberculosis (Latent or Active) Infection* Among Refugees By Region Of Origin, Minnesota, 2015


N=2,137 screened





Hepatitis B* infection Among Refugees by Region of Origin, Minnesota, 2015


N=2,084 screened





Intestinal Parasitic Infection* Among Refugees by Region of Origin, Minnesota, 2015

N=1,909 screened



*At least parasitic infection found via stool or serology (excluding nonpathogenic)



Health Status of New Refugees, Minnesota Immunization Status, 2004 – 2015

