

Adolescent Mental Health and/or Depression Screening Measure Specifications

2024 Measurement Year

0 001			
Summary of Changes	NONE		
Description	The percentage of patients 12-17 years of age who were screened for mental health and/or depression		
Measurement Period	FINAL 2024 MY: January 1, 2024 through December 31, 2024		
	PRELIMINARY 2025 MY: January 1, 2025 through December 31, 2025		
Eligible Population	Eligible Specialties	Family Medicine, Internal Medicine, Pediatric/Adolescent Medicine	
	Eligible Providers	Medical Doctor (MD), Doctor of Osteopathy (DO), Physician Assistant (PA), Advanced Practice Registered Nurses (APRN)	
	Ages	12 years or older at the start of the measurement period AND less than 18 years at the end of the measurement period	
	Event	At least one office well child visit or eligible telehealth encounter (Adolescent Well Child Visit Value Set) performed or supervised by an eligible provider in an eligible specialty during the measurement period	
Denominator	The eligible population		
Numerator	The number of patients in the denominator who completed ANY of the following mental health and/or depression screening tools during the measurement period AND have the screening tool result documented in the medical record: • Patient Health Questionnaire – 9 item version (PHQ-9) • PHQ-9M Modified for Teens and Adolescents • Kutcher Depression Scale (KADS) • Beck Depression Inventory II (BDI-II)		
	Beck Depression Inventory Fast Screen (BDI-FS)		
	 Child Depression Inventory (CDI) Child Depression Inventory II (CDI-2) 		
	Patient Health Qu	sestionnaire – 2 item version (PHQ-2)	
	Pediatric Sympton	n Checklist – 17 item version (PSC-17) - parent version	
	Pediatric Sympton	n Checklist – 35 item (PSC-35) - parent version	
	Pediatric Sympton	n Checklist – 35 item Youth Self-Report (PSC Y-SR)	
	Global Appraisal of substance abuse (0)	of Individual Needs screens for mental health and GAIN-SS)	



Adolescent Mental Health and/or Depression Screening Measure Specifications

2024 Measurement Year

Required Exclusions	The following exclusions must be applied to the eligible population:	
	Patient had a diagnosis of schizophrenia (Schizophrenia or Psychotic Disorders Value Set) any time prior to the end of the measurement period	
	Patient had a diagnosis of bipolar disorder (<i>Bipolar Disorder</i> Value Set) any time prior to the end of the measurement period	
	Patient had an active diagnosis of depression or dysthymia (Major Depression or Dysthymia Value Set) any time during the measurement period	
	Patient had a diagnosis of personality disorder (<i>Personality Disorder</i> – <i>Emotionally Labile</i> Value Set) any time prior to the end of the measurement period	
	Patient had moderate, severe, or profound intellectual disabilities (<i>Intellectual Disability</i> Value Set) any time prior to the end of the measurement period	
Measure Scoring	Rate/Proportion	
Interpretation of Score	Higher score indicates better quality	
Measure Type	Process	