

Contents


Appendix H: Baseline Responses on Symptom Measures in the Patient Self-Evaluation.....	H-1
Standard 8 Symptom Responses at Baseline.....	H-1
ANXIETY	H-1
APPETITE LACK	H-2
DEPRESSION	H-2
DISTURBED SLEEP.....	H-3
FATIGUE.....	H-3
NAUSEA	H-4
PAIN.....	H-4
VOMITING	H-5
PEG Scale Composite Scores At Baseline.....	H-6

Appendix H: Baseline Responses on Symptom Measures in the Patient Self-Evaluation

Standard 8 Symptom Responses at Baseline

ANXIETY


Figure 1. Distribution of standard 8 anxiety responses (0-10 numerical rating scale; NRS) at baseline in intractable pain (IP) patients.



Patients falling to the right of the vertical, dotted red line (n = 1679) were patients experiencing moderate to severe anxiety at baseline.

APPETITE LACK


Figure 2. Distribution of standard 8 appetite lack responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 1154) were patients experiencing moderate to severe lack of appetite at baseline.

DEPRESSION


Figure 3. Distribution of standard 8 depression responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 1453) were patients experiencing moderate to severe depression at baseline.

DISTURBED SLEEP


Figure 4. Distribution of standard 8 disturbed sleep responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 1977) were patients experiencing moderate to severe sleep disturbance at baseline.

FATIGUE


Figure 5. Distribution of standard 8 fatigue responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 2036) were patients experiencing moderate to severe fatigue at baseline.

NAUSEA


Figure 6. Distribution of standard 8 nausea responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 1028) were patients experiencing moderate to severe nausea at baseline.

PAIN


Figure 7. Distribution of standard 8 pain responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 2165) were patients experiencing moderate to severe pain at baseline.

VOMITING


Figure 8. Distribution of standard 8 vomiting responses (0-10 NRS) at baseline in IP patients.



Patients falling to the right of the vertical, dotted red line (n = 437) were patients experiencing moderate to severe vomiting at baseline.

PEG Scale Composite Scores At Baseline

Figure 9. Distribution of PEG composite scores at baseline in IP patients.



For ease of display, composite scores were rounded down to the next whole number. Patients falling to the right of the vertical, dotted red line (n = 2129) were patients scoring 4 or above at baseline.